

Semiclassical analysis of the Nonequilibrium Local Polaron

A. Mitra, I. Aleiner and A. J. Millis

Department of Physics, Columbia University, 538 W. 120th St, NY, NY 10027

(Dated: February 2, 2008)

A resonant level strongly coupled to a local phonon under nonequilibrium conditions is investigated. The nonequilibrium Hartree-Fock approximation is shown to correspond to approximating the steady state density matrix by delta functions at field values to which the local dynamics relaxes in a semiclassical limit. If multiple solutions exist, all are shown to make nonvanishing contributions to physical quantities: multistability does not exist. Nonequilibrium effects are shown to produce decoherence, causing the standard expansions to converge and preventing the formation of a polaron feature in the spectral function. The formalism also applies to the nonequilibrium Kondo problem.

PACS numbers: 73.23.-b, 05.30.-d, 71.10.-w, 71.38.-k

The quantum mechanics of nonequilibrium systems is a subject of fundamental importance and of great current interest, for example in the context of prospective 'single molecule devices'¹. In equilibrium problems, non-perturbative analysis based on solutions of Hartree-Fock equations (which may be understood as saddle points of functional integrals) has led to important insights. In nonequilibrium problems, mean field equations may be formulated^{2,3,4} and indeed exhibit a richer structure than in the corresponding equilibrium problems, but it has not been clear how to select the relevant solutions or to systematically compute corrections.

In this paper we investigate a simple model which indicates a resolution to these issues. We find that the relevant solutions are selected by the steady state density matrix, which in a semiclassical, weakly nonequilibrium limit is found to become very sharply peaked at field values corresponding to local minima of a 'pseudoenergy' which we define. The formalism also shows that if multiple minima exist, all contribute, with weights varying smoothly as parameters change and shows how departures from equilibrium lead to a decoherence which suppresses characteristically quantal effects such as the formation of a polaron resonance.

We consider a single level which may be occupied by 0 or 1 spinless electrons (creation operator $d^\dagger$) and is coupled to two leads $j = L, R$ (creation operators $a_{jk}^\dagger$). The leads are assumed to be reservoirs specified by chemical potentials μ_j and inverse temperature β_j (which we typically set to $\beta_j = \infty$). The electrons interact with an oscillator (dimensionless displacement coordinate q and momentum p) of mass M_{ph} and energy $U(q)$. This is a local version of the familiar 'polaron problem'⁵ and captures an important aspect of the physics of prospective single-molecule devices^{6,7}. The Hamiltonian is $H = H_{ce} + H_{tun} + H_{dot} + H_{ph} + H_I$; with

$$H_{ce} = \sum_{k,j} \epsilon_{k,a} a_{k,j}^\dagger a_{k,j}; \quad H_{dot} = \epsilon_0 d^\dagger d; \quad (1)$$

$$H_{tun} = \sum_{k,j} t_{k,j} (d^\dagger a_{k,j} + h.c) \quad (2)$$

$$H_I = \lambda q d^\dagger d; \quad H_{ph} = \frac{p^2}{2M_{ph}} + U(q) \quad (3)$$

At $M_{ph} = \infty$ the model is analytically solvable, with properties determined by the Green functions

$$\mathbf{g}_q(t-t') = \begin{pmatrix} g_{qR}(t-t') & g_{qK}(t-t') \\ 0 & g_{qA}(t-t') \end{pmatrix} \quad (4)$$

where $g(t) = \int \frac{d\omega}{2\pi} e^{-i\omega t} g(\omega)$ with

$$g_{qR}(\omega) = \frac{1}{\omega - \epsilon_0 - \lambda q - \Sigma(\omega + i\delta)} = g_{qA}(\omega)^* \quad (5)$$

$$g_{qK}(\omega) = 2i \sum_{j=L,R} [a_j(\omega) \tanh(\beta_j(\omega - \mu_j))], \quad (6)$$

and $\Sigma = \Sigma_L + \Sigma_R$, $\Sigma_j(z) = \sum_p \frac{t_{p,j}^2}{z - \epsilon_{pa}}$ and $a_j = Im \Sigma_j(\omega - i\delta) / ((\omega - \epsilon_0 - \lambda q - Re \Sigma(\omega))^2 + Im \Sigma^2(\omega))$. It is also useful to define $b = Reg_R$ and the 'pseudoenergy'

$$\Phi(q) = U(q) + \int_0^q dq' \lambda q' \int \frac{d\omega}{2\pi i} \frac{g_{q'K} - g_{q'R} + g_{q'A}}{2} \quad (7)$$

which in equilibrium becomes the q -dependent part of the total energy. Fig. 1 shows a possible form of $\Phi(q)$.

The equilibrium physics of this model is very well understood, and is conveniently viewed as a functional integral over trajectories $q(t)$. In the limits $M_{ph}, \beta \rightarrow \infty$ the integral is dominated by those paths for which q takes the definite value q_a minimizing the energy. At finite M_{ph} , other paths become important: two crucial classes are the 'gaussian fluctuation' paths involving small excursions (characteristic frequency ω_a) from the minimum and, if $\Phi_{eq}(q)$ has the form shown in Fig 1 and the barrier height H is sufficiently large, tunnelling (instanton) processes during which q goes rapidly from the vicinity of the global minimum (E_1 in the notations of Fig 1) to the higher energy minimum (here E_2), spends a time of order $\Delta E^{-1} \equiv (E_2 - E_1)^{-1}$ near the higher minimum, and then returns to the vicinity of the global minimum. If H is sufficiently large, then tunnelling processes are rare, but when $E_2 \approx E_1$, the time spent in the higher minimum becomes longer than the interval




FIG. 1: Assumed dependence of 'pseudoenergy' (Eq. 7) on oscillator coordinate q .

between tunnelling events and the dilute instanton approximation breaks down, signalling the formation of a polaron resonance in the density of states. In this paper we investigate the changes occurring when the system is driven out of equilibrium by application of a chemical potential difference $\Delta\mu$ between the two leads. We present a general discussion but focus most attention on the nonequilibrium polaron limit $H \gg \omega_a \gg \Delta\mu$; with $\beta_j = \infty$ but $\Delta E/\Delta\mu$ arbitrary. In our actual calculations we also assume that departures from equilibrium are small enough that we may neglect density-of-states variations: $(\Delta\mu)\partial \ln a / \partial \omega < 1$.

To analyse the out of equilibrium behavior we use the standard^{8,9} extension of Feynman diagrammatics to the *Keldysh contour* which consists of a time-ordered (-) branch extending from $t = -\infty$ to $t = +\infty$, followed by an anti-time-ordered (+) branch extending from $+\infty$ back to $-\infty$. The Keldysh diagrammatics may be obtained by functional differentiation, with respect to source terms $\eta_a(t)$, of a generating functional $W[\{\eta_a\}]$ which can be formulated² as a coherent state path integral:

$$W[\{\eta_a\}] = \int dq_+ dq_- \hat{\rho}(q_+, q_-) \int \mathcal{D}q(t) \hat{S}_K[-\infty_+, -\infty_-] \quad (8)$$

Here the functional integral $\int \mathcal{D}q(t)$ is over all paths beginning at q_- at $t = -\infty$ on the time ordered contour and ending at q_+ at $t = -\infty$ on the antitime ordered contour, while $\hat{S}_K[t_1, t_2; \eta_a(t)] = T_K e^{i \int_{t_2}^{t_1} dt a(t) \hat{H}[\eta_a(t)]}$ the time evolution operator on the Keldysh contour and $a(t) = \pm 1$ according to whether the time is on - or + branch. The contributions of paths with endpoints q_+, q_- are weighted by the appropriate element of the steady state density matrix ρ which is the long time limit of an equation which may be expressed in path integral language as

$$\hat{\rho}(q_+, q_-; t; \eta) = \int dq'_+ dq'_- \int \mathcal{D}'q'(t') \hat{S}_K[t_+, t_0+] \hat{\rho}(q'_+, q'_-; t_0) \hat{S}_K[t_0-, t_-] \quad (9)$$

Here the $\int \mathcal{D}'q'(t')$ is over all paths which begin at q'_- on the lower contour at time t_0 and end at q'_- time t on the lower contour, along with the time reversed paths which begin at $q'_+(t)$ on the upper contour and return to q_+ at time t_0 . We require the long time behavior after transients have decayed. We do not find non-steady long time limits such as limit cycles or chaos: ρ in Eq 8 is the time independent solution of Eq 9.

Because eqs 1,2,3 involve a finite system coupled to two infinite reservoirs, the trace over electron operators may be performed⁷. The combination of time evolution operators needed in eqs. 8,9 may be written in the interaction representation as $\mathcal{R}(t_2, t_1; \{q(t)\}) \equiv \langle T_K \text{Exp}[-i \int_{t_1}^{t_2} dt (\lambda q_c(n_- - n_+) + \lambda_q q_q(n_- + n_+)) \rangle$ where $q_c(t) = \frac{q_-(t) + q_+(t)}{2}$, $q_q(t) = \frac{q_-(t) - q_+(t)}{2}$, $\langle \rangle$ denotes expectation value in the reservoir defined by the leads (note this trace depends on the entire trajectory $q(t)$) and $n_{+/-} = d^\dagger d$ on the indicated contour. In the physical problem the quantum field coupling constant $\lambda_q = \lambda$. Differentiation with respect to λ_q and application of the usual linked cluster arguments leads to

$$\mathcal{R}(t_2, t_1; \{q_c(t), q_q(t)\}) = e^{-\int_{t_1}^{t_2} dt (C(t) + \mathcal{L}_{ph+S})} \quad (10)$$

Here $\mathcal{L}_{ph+S}$ is the phonon action $M_{ph}(\partial_t q_q \partial_t q_c) - U(\frac{q_+ + q_-}{2}) - U(\frac{q_+ - q_-}{2})$ supplemented by source terms,

$$C(t) = q_q(t) \int_0^\lambda d\lambda_q \text{Tr} [\sigma_x \mathbf{G}_{\lambda_q}(t, t)] \quad (11)$$

and $\mathbf{G}_{g'}$ solves

$$\mathbf{G}_{\lambda_q} = \mathbf{g} + \mathbf{g} * \mathbf{v} * \mathbf{G}_{\lambda_q} \quad (12)$$

with $\mathbf{v}(t) = (\lambda q_c(t) \mathbf{1} + \lambda_q q_q(t) \sigma_x)$. We have verified that an expansion in powers of λ about the weak coupling limit $\lambda = 0$ reproduces results obtained⁷ by standard Keldysh diagrammatic analysis.

We now turn to the semiclassical analysis. In the large M_{ph} limit, one expects stationary (time-independent) paths to dominate the physics. However, in the Keldysh formalism, stationary paths have $q_q = 0$, so in the absence of source terms the time evolution operator in Eq 8 is unity for all stationary paths, providing no basis for selection. Instead, the important paths are selected by the density matrix; i.e. from the solution of Eq 9.

To understand the dynamics implied by Eq 9 we first assume that $\rho(q, q')$ is strongly peaked near $q = q' = q_a$. We may then expand $\Phi = E_a + \frac{M_{ph}\omega_a^2}{2} (q - q_a)^2$ and analyse Eq 9 by the usual perturbative methods⁷. In the large mass, weakly nonequilibrium limit we find relaxation, with a rate of order $\lambda^2 \omega_a (a_L + a_R)^2$ to a sum of functions sharply peaked about the values q_a which minimize the pseudoenergy.

$$\rho(q, q') \approx \sum_a \rho_a r_a(q, q') \quad (13)$$

To leading order in the electron-phonon coupling λ we find $r_a(q, q') = \int \frac{dp}{2\pi} e^{ip(q-q')} \text{Exp}[-(\frac{p^2}{2M_{ph}} +$

$\frac{M_{ph}\omega_a^2(\frac{q+q'}{2}-q_a)^2}{2})\frac{2\tanh[\frac{\omega_a}{2T_{eff}}]}{\omega_a}]$ with $T_{eff} = \Delta\mu\frac{a_L a_R}{(a_L+a_R)^2}$, peaked about q_a with a width arising from quantum fluctuations (finite M_{ph}) and from departures from equilibrium (which act as an effective temperature).

Fixing the ρ_a requires consideration of the exponentially small processes neglected above. Within the present formalism we find two: quantal (finite M_{ph}) effects which lead to tunnelling through the barrier connecting the minima, and diffusion (finite $\Delta\mu$) effects which produce motion along the pseudoenergy surface. Diffusion may be analysed using the analogy between $\Delta\mu$ and temperature mentioned above; it leads¹² to motion between minima with rate constant R_{diff} of order $\ln R_{diff} \approx \frac{H}{\Delta\mu}$ if $\Delta\mu \gg \omega_a$ and to a much smaller rate in the opposite limit. The standard equilibrium estimate of the rate due to tunnelling processes gives $\ln R_{tun} \approx -2\sqrt{\frac{HK_a(q_1-q_2)^2}{\omega_a}} \sim \frac{H}{\omega_a}$; corrections become important only for $\Delta\mu \sim \omega_a$. Thus we expect roughly that for $\Delta\mu > \omega_a$, the diffusion process dominates.




FIG. 2: Instanton processes entering equation of motion of density matrix.

We now consider in more detail the interesting quantum limit $\Delta\mu < \omega_a$ for large barrier height H . In this limit one would like to restrict attention to paths for which $q(t)$ is almost always near one of the minima, with occasional tunnelling events in which q shifts from one minimum to another. The tunnelling amplitude R_{tun} is exponentially small and (for small enough $\Delta\mu$) is well approximated by its equilibrium value. However, in the real time path integral the smallness of R_{tun} arises as the nearly complete cancellation of a sum of many paths with large but oscillating amplitudes. We argue that for small $\Delta\mu$ the result of performing this complicated sum (which we do not treat directly) may be represented by the 'instanton' (kink) vertices shown in Fig 2 with amplitude R_{tun} . A kink or antikink comes with a factor i and a sign $\sigma = \pm 1$ determined by the contour it is on. We find (see below) that the action of a single kink or antikink is infinite, so each kink must be followed by an

antikink. The path integral Eq 9 thus becomes

$$\rho_a(t) = \rho_a(t_0) - R_{tun}^2 \int_{t_0}^t dt_f dt_i (\mathcal{R}(t_f, t_i; \{q_{ca} \rightarrow q_{ca}\}) \rho_a(t_0) - \mathcal{R}(t_f, t_i; \{q_{ca} \rightarrow q_{cb}\}) \rho_b(t_0)) + \mathcal{O}R^4 \quad (14)$$

where the first term is the sum of the processes shown in panels *a* and *b* in Fig. 2 the second term is the sum of the processes shown in *c* and *d*, and we have denoted explicitly only the beginning and ending values of the classical component of the field.

Differentiating Eq 14 with respect to time yields

$$\frac{d\rho_a(t)}{dt} = -\Gamma_{a \rightarrow b} \rho_a(t) + \Gamma_{b \rightarrow a} \rho_b(t) \quad (15)$$

where to order R_{tun}^2 the scattering rates are

$$\Gamma_{a \rightarrow a'} = R_{tun}^2 Re \int_0^\infty dt \mathcal{R}(t, 0; \{q_{ca} \rightarrow q_{ca'}\}) \quad (16)$$

The integral in Eq 16 yields a quantity T_0 with dimension of time; we may neglect higher order terms if $R_{tun}T_0 \ll 1$. To evaluate Eq 16 it suffices to approximate the potential in Eq 12 by the 'telegraph' form shown in Fig 2; the equations may then be solved by standard methods. In equilibrium a crucial role is played by the phase shift $\delta = \tan^{-1} \left(\frac{(a_L+a_R)V}{1-bV} \right)$ ¹⁰ with $V = \lambda(q_b - q_a)$ and $a_{L,R}$ and b given by Eqs 5,6 at $q = q_a$. As shown by Ng¹¹, out of equilibrium, the behavior is described by complex phase shifts $\delta_{L,R}$ with

$$\delta_{L,R} = \tan^{-1} \left(\frac{a_{L,R}V}{1-bV - ia_{R,L}V \text{sgn}(\mu_{L,R} - \mu_{R,L})} \right) \quad (17)$$

Note $\delta_L + \delta_R = \delta$. We find

$$\ln \mathcal{R} = (C_{eq}(\bar{\mu}) - i\delta E_{neq} + \delta C_{dec} + \delta C_{orth}) t_{fi} \quad (18)$$

Here $\delta E_{neq} = (\delta'_L - \delta'_R) \Delta\mu$ is the change in electronic energy arising from the imbalance in chemical potential. $C_{eq}(\bar{\mu}) - i\delta E_{neq}$ is the equilibrium result evaluated at the mean chemical potential plus the nonequilibrium energy correction; it oscillates at frequency ΔE , providing convergence of the integral in Eq 14 when $\Delta E \neq 0$. The imaginary parts δ'' of the phase shifts give rise to a time decay ('decoherence') which is the manifestation, in the present formalism, of the decoherence introduced on semiphenomenological grounds by Rosch et. al.¹³. and ensures that out of equilibrium the integral in Eq 14 converges even when $\Delta E + \Delta E_{neq} = 0$. We find

$$\delta C_{dec} = -\frac{(\delta''_L - \delta''_R)}{2\pi} \Delta\mu \phi_2(\Delta\mu t_{fi}) \quad (19)$$

At short times $\Delta\mu t_{fi} \ll 1$, $\delta C_{dec} t_{fi} \sim -(\Delta\mu t_{fi})^2$; at long times $\delta C_{dec} t_{fi} \sim -|\Delta\mu t_{fi}|$. At intermediate times an analytical solution is not available and ϕ_2 must




FIG. 3: *Main panel*: Variation of current with chemical potential difference $\Delta\mu$ at fixed pseudoenergy difference $E_1 - E_2$ calculated for parameters $\lambda q_1 = 4\Sigma''$ (resonance well above fermi level) and $\lambda q_2 = -\Sigma''$ (resonance slightly below fermi level) and equal coupling to left and right leads. *Upper Inset*: Spectral functions corresponding to equilibrium state (solid curve—only minimum "1" occupied) and strongly nonequilibrium state (dashed curve; incoherent superposition of the G 's corresponding to the two minima). *Lower Inset*: Variation with pseudoenergy difference of density matrix element corresponding to states near pseudoenergy minimum 1, computed for parameters used in main panel.

be computed perturbatively or numerically. We find $\phi_2(x) = \left(\frac{2}{\pi}\right) \left(Si(x) - \frac{1-\cos(x)}{x}\right)$.

δC_{orth} expresses the change in orthogonality effects arising because at nonvanishing $\Delta\mu$ there is destructive interference between the left and right leads¹¹:

$$\delta C_{orth} t_{fi} = \frac{2Re\delta_L\delta_R}{\pi^2} \psi_1(\Delta\mu t) + \frac{2iIm\delta_L\delta_R}{\pi^2} \psi_2(\Delta\mu t) \quad (20)$$

with $\psi_1(x) = \gamma_e - Ci(x) + \ln(x)$ and $\psi_2(x) = \frac{2}{\pi} \int_0^1 du \sin(ux) \frac{[(1-u)\ln(1-u)+u\ln u]}{u^2}$.

We have used the perturbative expressions for the

crossover functions to evaluate Γ (Eq 16). In equilibrium and at $\beta_j = \infty$ we find that only scattering from the higher energy to the lower energy extremum occurs (the 'up-scattering rate' vanishes); under nonequilibrium conditions an upscattering rate appears: of order $\left(\frac{\Delta\mu}{\Delta E}\right)^{\left(\frac{\Delta E}{\Delta\mu}\right)}$ if $\frac{|\Delta E|}{\Delta\mu} \gg 1$ but of the order of the down-scattering rate as $\frac{|\Delta E|}{\Delta\mu} \rightarrow 0$. The change in the weight ρ_1 for one of the two minima as $\Delta\mu/\Delta E$ is varied is shown in the lower inset of Fig 3. A similar result was found by Parcollet and Hooley in a 'pseudofermion' diagrammatic study of the nonequilibrium Kondo problem¹⁴.

The semiclassical Greens function follows as:

$$G_R(\omega) = \sum_a \frac{\rho_a((\Delta\mu))}{\omega - \epsilon_0 - gq_a - \Sigma_R(\omega)} \quad (21)$$

The main panel of Fig 3 shows the current computed in the standard way by inserting the Green function given in Eq. 21 into Eq 28 of Ref⁷. The inset shows the evolution of the spectral density $A(\omega) = ImG_R(\omega)$.

The minimum condition $\partial\Phi/\partial q = 0$ is equivalent to the Hartree-Fock equations discussed in recent literature^{3,4}, but in these works it is assumed that at each set of parameter values, only one of the minima is occupied ($\rho_a = 0$ or 1); preparation conditions are argued^{3,4} to determine the state of the system, leading to multistability and switching, in contradiction to the results shown in Fig 3. Although the bistability discussed by^{3,4} does not occur, the slow dynamics governing equilibration between the minima will lead to 'telegraph' noise in the current.

Hamann¹⁵ shows that interacting electron problems such as the Anderson model lead, after Hubbard-Stratonovich transformation, to a model very like that studied here, but with the role of the phonon field played by the spin part of the decoupling field. Our analysis carries over directly to these problems, providing a different derivation¹⁶ of the generally accepted results^{13,14} that nonequilibrium effects suppress the formation of the Kondo resonance, and that the nonequilibrium Kondo effect is fundamentally a weak coupling problem.

Acknowledgements: We thank the Columbia University Nanocenter and nsf-dmr0338376 for support, and L.Glazman for very helpful conversations.

¹ See, e.g. A.W. Ghosh, P.S. Damle, S. Datta, and A. Nitzan, MRS Bulletin **29**, 391 (2004).

² For a review, see, e.g. A. Kamenev, unpublished (cond-mat/0109316).

³ A. S. Alexandrov, A. M. Bratkovsky, and R. Stanley Williams Phys. Rev. **B67**, 075301 (2003).

⁴ A. Komnik and A. Gogolin, Phys. Rev. **B69** 153102 (2004) and cond-mat/0207513.

⁵ G. Mahan, *Many Particle Physics*, Chapter 4, 2nd ed. (Plenum, New York, 1990).

⁶ H. Park, J. Park, A.K.L. Lim, E.H. Anderson, A.P. Alivisatos, and P.L. McEuen, Nature **407**, 57 (2000).

⁷ A. Mitra, I. Aleiner and A. J. Millis, Phys. Rev. **B69**, 245302 (2004).

⁸ L.V. Keldysh, Zh. Éksp. Teor. Fiz. **47**, 1945 (1964) [Sov. Phys. JETP **20**, 1018 (1965)]

⁹ J. Rammer, *Quantum Transport Theory*, (Perseus: Reading MA, 1998).

¹⁰ P. Nozières and C. T. De Dominicis, Phys. Rev. **178**, 1097-1107 (1969).

- ¹¹ Tai-Kai Ng, Phys. Rev. **B54**, 5814 (1996).
- ¹² H. A. Kramers; Physica **7** 284 (1940).
- ¹³ A. Rosch, J. Paaske, J. Kroha, and P. Wölfle, Phys. Rev. Lett. **90** 076804/1-4 (2003).
- ¹⁴ O. Parcollet and C. Hooley, Phys. Rev. **B66**, 085315 (2002).
- ¹⁵ D. R. Hamann, Phys. Rev. **B2**, 1373 (1970).
- ¹⁶ A. Mitra and A. J. Millis, in preparation.